

Cross Section Evaluation by Spinor Integration I: The massless case in 4D

Bo Feng^{◇†}, Rijun Huang^{*}, Yin Jia^{*}, Mingxing Luo^{*}, Honghui Wang^{*}

[◇]*Center of Mathematical Science, Zhejiang University, Hangzhou 310027, China*

^{*}*Zhejiang Institute of Modern Physics, Physics Department, Zhejiang University, Hangzhou 310027, China*

[†]*Division of Applied Mathematica and Theoretical Physics,
China Institute for Advanced Study, Central University of Finance
and Economics, Beijing, 100081, China*

ABSTRACT: To get the total cross section of one interaction from its amplitude $\mathcal{M}$, one needs to integrate $|\mathcal{M}|^2$ over phase spaces of all out-going particles. Starting from this paper, we will propose a new method to perform such integrations, which is inspired by the reduced phase space integration of one-loop unitarity cut developed in the last few years. The new method reduces one constrained three-dimension momentum space integration to an one-dimensional integration, plus one possible Feynman parameter integration. There is no need to specify a reference framework in our calculation, since every step is manifestly Lorentz invariant by the new method. The current paper is the first paper of a series for the new method. Here we have exclusively focused on massless particles in 4D. There is no need to carve out a complicated integration region in the phase space for this particular simple case because the integration region is always simply $[0, 1]$.

KEYWORDS: Unitarity Cut, Cross Section.

Contents

1. Introduction	2
1.1 The reduced phase space integration of unitarity cut	4
2. Spinor method for the physical phase space	7
2.1 The phase space integration with three out-going particles	7
2.2 Arbitrary number of out-going particles and the recursive method	10
3. Example one: two out-going particles	11
3.1 The pure phase space integration	11
3.2 The example with $f = 2L_1 \cdot L_2$	12
3.3 The example with $f = 1/(2P_1 \cdot L_2 + m^2)$	13
3.4 The divergent behavior	13
4. Example two: three out-going particles	15
4.1 The pure phase space integration	15
4.2 The example with $f = (L_1 + L_3)^2$	16
4.3 The example with $f = s^2 t$	16
5. Example three: four out-going particles	17
5.1 The pure phase space integration	17
5.2 Two examples of non-trivial f	17
6. Conclusion	19
A. Using momentum method to integrate f_2 for four out-going particles	20
B. The spinor integration as integration in complex plane	22
B.1 Rewriting of spinor integration	22
B.2 More examples	24

1. Introduction

In the last few years there were great progresses in the evaluation of one-loop amplitudes for general field theory (see [1] and references within). One of such progresses is the unitarity cut method, which was initiated in [2, 3] and then pushed by Witten’s “twistor program” [4]. One key achievement along this line is the reduced unitarity phase space integration using “holomorphic anomaly” [5]. More accurately, the reduced unitarity phase space integration is given by¹

$$\int d^4L \delta^+(L^2) \delta^+((L - K)^2) G(L) . \quad (1.1)$$

With the first delta-function $\delta^+(L^2)$ we can reduce the measure $\int d^4L \delta^+(L^2)$ into the integration with variables $t, \lambda, \tilde{\lambda}$, where t is a affine variable and $\lambda, \tilde{\lambda}$ are spinor variables. With the second delta-function $\delta^+((L - K)^2)$, we can integrate t out and get

$$\int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] f(\lambda, \tilde{\lambda}) . \quad (1.2)$$

Before the work of [5], the evaluation of remaining two-dimensional integration is a very hard task and has blocked practical applications of unitarity cut method. From [5] is that people realized that the remaining integrations over $\lambda, \tilde{\lambda}$ can be obtained by reading out residues of corresponding poles. In another word, there is no need for integrations and everything is just algebraic manipulation. This method is usually referred to as “Spinor Integration Method” or the spinor method for short.

Clearly, the success of the spinor integration method is due to the presence of two delta-functions. Using this technique, we are able to perform any unitarity phase space integration at one-loop in pure 4 dimension[6, 7, 8]. Originally, spinor is tightly related to null momentum in pure 4D. However, for practical applications, it will be useful to generalize it to general D-dimension as well as massive particles. This goal has been achieved late in [9, 10].

To get the cross section, we need to integrate the physical phase space² of all (or some) out-going particles,

$$c \int \prod_f \frac{d^3p_f}{2E_f} |\mathcal{M}|^2 \delta^4(P_{in} - P_{out}) \quad (1.3)$$

where $\delta^4(P_{in} - P_{out})$ is the energy-momentum conservation condition and c is a function depending on the in-coming particles (for example, $c^{-1} = 4E_A E_B |v_A - v_B|$ in $2 \rightarrow n$ progresses). For simplicity, we omit c in this paper.

¹The $\delta^+()$ means the integrated region with $l_0 \geq 0$.

²To make things clear, we use unitarity phase space and physical phase space to distinguish these two cases.

Usually, integrations as such are difficult to perform, especially when there are many out-going particles. For LHC experiments, channels with four or even five out-going particles play important role in search of new physics (see, for example, [11] and references in this paper). Aiming at this task, we will try to do the physical phase space integration using the spinor integration method.

The advantage of the spinor method is that it reduces the constrained three-dimension momentum space integration to an one-dimensional integration, plus one possible Feynman parameter integration. All remaining integrations are scalar type, i.e., the integrand is manifestly Lorentz invariant, so there is no need to specify a reference framework in our calculation. Furthermore the integration region in the phase space also becomes simpler.

The current paper is the first of a series of work we will take to complete our new method. In this paper, we will focus on massless particles in 4D. For this case, with familiar infrared/collinear divergences for massless out-going particles, there will be infinity after the phase space integration. To get sensible physical quantities, some regularization is needed. Due to this difficulty, the result in this paper is still far from the practical applications. It is the basis for all late work. We will deal with massive particles in the second paper and general D-dimension integration in the third part.

For massless particles in pure 4D as focused on in this paper, some simplifications happen. For example, the integration region in phase space will always be $[0, 1]$ in our method. This will be modified to be nontrivial functions of mass and energy when the particles are massive as to be presented in second paper.

The main aim of this paper is to laid out the framework of our new method, so most examples in this paper are not for real cross sections. These examples is to demonstrate the salient character of spinor integration, such as frame independence and simple integration region.

The outline of this paper is the following. In section 2, we transfer the physical phase space integrations into the form of spinor integrations. We start with the case when there are two and three outgoing particles, then generalize to cases with $n \geq 3$ recursively. The recursive feature (not recursion relation) is one of advantages of the spinor integration method.

In the following three sections, our method has been demonstrated with simple examples with two, three and four outgoing particles, respectively. There will also a brief discussion on the IR/collinear divergence problem related to massless particles in section 3.

In section 6, we summarize our results along with general discussions and remarks.

There are two appendixes. In Appendix A the evaluation of pure four particle phase space directly using momentum components has been given as to compare with spinor integration method presented in main text. In Appendix B we have explained the unfamiliar spinor integration method from the point of view of integration in complex plane. From this point, the correctness and the power of this method becomes obvious.




Figure 1: Representation of the cut integral. Left and right legs are on-shell. Internal lines represent the cut propagators.

1.1 The reduced phase space integration of unitarity cut

As we have emphasized above, the key gradient of spinor method is the existence of two delta-functions. There is one situation where these two delta-functions arise naturally: it is the reduced phase space integration of one-loop unitarity cut. In this subsection, we review how to perform the phase space integration of one-loop unitarity cut by spinor method.

Unitarity Cut: Consider the unitarity cut in the $(i, i+1, \dots, j-1, j)$ -channel, as shown in fig. 1. The cut integral is

$$C_{i,i+1,\dots,j-1,j} = \int d\mu A^{\text{tree}}(l_1, i, i+1, \dots, j-1, j, l_2) A^{\text{tree}}((-l_2), j+1, j+2, \dots, i-2, i-1, (-l_1)), \quad (1.4)$$

where $d\mu = d^4l_1 d^4l_2 \delta^{(+)}(l_1^2) \delta^{(+)}(l_2^2) \delta^4(l_1 + l_2 - P_{ij})$ is the Lorentz invariant phase space measure of two light-like vectors (l_1, l_2) constrained by the energy-momentum conservation, and $P_{ij} = p_i + \dots + p_j$. Notice that the integrand is similar to $|\mathcal{M}|^2$ in eq (1.3). So we can make full use of this point in our calculations.

Using the $\delta^4(\)$ function to perform the l_2 integration gives

$$C_{i,i+1,\dots,j-1,j} = \int d^4l_1 \delta^{(+)}(l_1^2) \delta^{(+)}((l_1 - P_{ij})^2) \times A^{\text{tree}}(l_1, i, i+1, \dots, j-1, j, l_2) A^{\text{tree}}(-l_2, j+1, j+2, \dots, i-2, i-1, -l_1) \quad (1.5)$$

Note that the Lorentz invariant measure of a null vector l can be represented as a measure over $\mathbb{R}^+ \times \mathbb{CP}^1 \times \mathbb{CP}^1$. The contour of integration is a certain diagonal $\mathbb{CP}^1$. Explicitly, one writes

$l_{a\dot{a}} = t\lambda_a\tilde{\lambda}_{\dot{a}}$, and then³

$$\int d^4l \delta^{(+)}(l^2)(\bullet) \sim \int_0^\infty t dt \int_{\tilde{\lambda}=\bar{\lambda}} \langle \lambda d\lambda \rangle [d\tilde{\lambda} \tilde{\lambda}](\bullet), \quad (1.6)$$

where $(\bullet)$ represents a generic integrand up to an overall numerical factor. So we have⁴

$$C_{i,\dots,j} \sim \int_0^\infty t dt \int (-) \langle \lambda d\lambda \rangle [\tilde{\lambda} d\tilde{\lambda}] \delta^{(+)} \left(P_{ij}^2 - t \left\langle \lambda | P_{ij} | \tilde{\lambda} \right\rangle \right) G(\lambda, \tilde{\lambda}, t). \quad (1.7)$$

$G(\lambda, \tilde{\lambda}, t)$ arises from the product of the two tree-level amplitudes in (1.4). In order to get $G(\lambda, \tilde{\lambda}, t)$ in actual calculations, we have to write expressions of the form $\langle \bullet, l_2 \rangle$ or $[\bullet, l_2]$ in terms of $l = l_1$ (l_1 is always substituted by l when there is no possibility of confusion). A systematic way of doing this is as follows:

$$\langle \bullet, l_2 \rangle = \frac{\langle \bullet, l_2 \rangle [l_2 l_1]}{[l_2 l_1]} = \frac{\langle \bullet | l_2 | l_1 \rangle}{[l_2 l_1]} = \frac{\langle \bullet | P_{ij} | l_1 \rangle}{[l_2 l_1]}. \quad (1.8)$$

A similar identity is valid for $[\bullet, l_2]$. The factors $[l_2 l_1]$ and $\langle l_2 l_1 \rangle$ all pair up in the end allowing for the use of the vector form of l_2 . This is because the product of amplitudes must be invariant under the scaling $z\lambda_{l_2}$ and $z^{-1}\tilde{\lambda}_{l_2}$. For (1.7), integrating over t yields

$$C_{i,\dots,j} \sim P_{ij}^2 \int \frac{\langle \lambda d\lambda \rangle [\tilde{\lambda} d\tilde{\lambda}]}{\left\langle \lambda | P_{ij} | \tilde{\lambda} \right\rangle^2} G \left(\lambda, \tilde{\lambda}, \frac{P_{ij}^2}{\left\langle \lambda | P_{ij} | \tilde{\lambda} \right\rangle} \right). \quad (1.9)$$

The requirement of degree zero in $\tilde{\lambda}$ implies that G in (1.9) can be written as a sum of terms in the form⁵

$$\frac{\prod_{i=1}^k \left\langle \lambda | R_i | \tilde{\lambda} \right\rangle}{\left\langle \lambda | P_{ij} | \tilde{\lambda} \right\rangle^a \prod_{j=1}^{k-a} \left\langle \lambda | Q_j | \tilde{\lambda} \right\rangle}, \quad (1.10)$$

where R_i, Q_j are functions of external momenta.

Canonical Splitting: In order to calculate the integral efficiently, we can reduce the integrand by separating the denominator factors with $\tilde{\lambda}$ as much as possible, at the cost of more terms. When

³Here we have used the QCD convention so measure is $\langle \lambda d\lambda \rangle [d\tilde{\lambda} \tilde{\lambda}]$ instead of the twistor convention $\langle \lambda d\lambda \rangle [\tilde{\lambda} d\tilde{\lambda}]$.

⁴Where we have written $(\ell_1 - P_{ij})^2 = P_{ij}^2 - t \left\langle \lambda | P_{ij} | \tilde{\lambda} \right\rangle$, thus we have used the QCD convention instead of the twistor convention.

⁵It is worth to emphasize that this general form is only true for the one-loop calculation. When we apply our method to cross section evaluation, we will have other forms of input. The way to deal with them generally is given in Appendix B.

there is a product $[a \tilde{\lambda}][b \tilde{\lambda}]$ in the denominator, multiply both numerator and denominator by $[a b]$. Applying Schouten's identity yields

$$[i j][k l] = [i k][j l] + [i l][k j] \quad (1.11)$$

in the numerator with another factor $[c \tilde{\lambda}]$ (which must exist by homogeneity when the degree of $\tilde{\lambda}$ in denominator is equal to or more than three). Thus we get two terms with $[a \tilde{\lambda}]$ or $[b \tilde{\lambda}]$ in the numerator, canceling one of the denominator factors. The result, in terms of $\tilde{\lambda}$, is a denominator of the form $\prod_r \langle \lambda | Q_r | \tilde{\lambda} \rangle [A \tilde{\lambda}]$ in every term. Similarly $\langle \lambda | Q_r | \tilde{\lambda} \rangle$ can be treated by writing $\langle \lambda | Q_r | \tilde{\lambda} \rangle = [\tilde{Q}_r \tilde{\lambda}]$, where $\tilde{\lambda}_{Q_r} = -(Q_r)_{a\dot{a}} \lambda_l^a$.

Using this procedure repeatedly and noticing the integrand has a degree of -2 in $\tilde{\lambda}$, we end up with two kinds of possible integrals:

$$\mathcal{I}_A = - \int \frac{\langle \lambda d\lambda \rangle [\tilde{\lambda} d\tilde{\lambda}] \prod_{i=1}^k \langle \lambda | R_i | \tilde{\lambda} \rangle}{\langle \lambda | P_{ij} | \tilde{\lambda} \rangle^{2+k}}; \quad \mathcal{I}_B = - \int \frac{\langle \lambda d\lambda \rangle [\tilde{\lambda} d\tilde{\lambda}]}{\langle \lambda | P_{ij} | \tilde{\lambda} \rangle \langle \lambda | Q_r | \tilde{\lambda} \rangle}. \quad (1.12)$$

$\mathcal{I}_B$ can be changed to

$$\mathcal{I}_B = - \int_0^1 dx \int \frac{\langle \lambda d\lambda \rangle [\tilde{\lambda} d\tilde{\lambda}]}{\langle \lambda | x P_{ij} + (1-x) Q_r | \tilde{\lambda} \rangle^2}, \quad (1.13)$$

which reduces to $\mathcal{I}_A$ up to a Feynman parametrization integration. On the other hand, $\mathcal{I}_A$ is just the $\prod_i y_i$ component of an auxiliary integration⁶

$$\mathcal{I}_{aux} = \int \frac{\langle \lambda d\lambda \rangle [\tilde{\lambda} d\tilde{\lambda}] \langle \lambda | R | \tilde{\lambda} \rangle^k}{\langle \lambda | P_{ij} | \tilde{\lambda} \rangle^{2+k}}, \quad R = \sum_i y_i R_i, \quad (1.14)$$

Now all integrations are reduced into the one given by $\mathcal{I}_{aux}$. It can in turn be written as a total derivative by using

$$\frac{[\tilde{\lambda} d\tilde{\lambda}][\eta \tilde{\lambda}]^n}{\langle \lambda | P | \tilde{\lambda} \rangle^{n+2}} = [d\tilde{\lambda} \partial_{\tilde{\lambda}}] \left(\frac{1}{(n+1) \langle \lambda | P | \eta \rangle} \frac{[\eta \tilde{\lambda}]^{n+1}}{\langle \lambda | P | \tilde{\lambda} \rangle^{n+1}} \right). \quad (1.15)$$

Thus, the evaluation of formula (1.14) is transformed into the reading of residues of

$$\frac{1}{(k+1) \langle \lambda | R | P_{ij} | \lambda \rangle} \frac{\langle \lambda | R | \tilde{\lambda} \rangle^{k+1}}{\langle \lambda | P_{ij} | \tilde{\lambda} \rangle^{k+1}}$$

⁶There are other ways to do it directly without using the auxiliary integration. See reference [7].

where we need to sum up two possible poles from the factor $\langle \lambda | R | P_{ij} | \lambda \rangle$. The way to sum up these two poles has been discussed in detail, for example, in eq (70), (74) of [12]. In fact, the result can be written down directly as

$$\frac{-[(-2P \cdot R + \sqrt{\Delta})^{k+1} - (-2P \cdot R - \sqrt{\Delta})^{k+1}]}{(k+1)\sqrt{\Delta}(2P^2)^{k+1}} \quad (1.16)$$

with $\Delta = (2P \cdot R)^2 - 4P^2 R^2$. It is easy to see that numerator must be of the form $\sum_a \sqrt{\Delta}^{2a+1} (-2P \cdot R)^{k-2a}$. The result is a rational function with the highest power of R to be k , as required.

Before we end this part, let us emphasize that above procedure works for general one-loop calculation, but when we try to do the cross section calculation, we may meet new kinds of singularities and we need to generalize above procedure. The generalization has been discussed in Appendix B in some details. *The basis idea is still to find formula like the one in (1.15) and then take the residues.*

2. Spinor method for the physical phase space

In this section we will establish the general framework for the physical phase space integration using spinor method. We will focus on massless particles in pure 4D. For massive particles and in general D-dimension, it will be discussed subsequent work.

Before going into detail, let us make a simple observation: for two out-going particles, the physical phase space integration is the same as the unitarity phase space integration, which can be done directly by the spinor integration method. Difficulties arise when $n \geq 3$. We will first deal with the case $n = 3$ and then generalize to arbitrary out-going particles.

Notice that we can rewrite

$$\int \frac{d^3 p_f}{2E_f} \sim \int d^4 p_f \delta^+(p_f^2), \quad (2.1)$$

Then when there are only two out-going particles, we have

$$\prod_{f=1,2} \int d^4 p_f \delta^+(p_f^2) \delta^4(P_{in} - p_1 - p_2) = \int d^4 p_1 \delta^+(p_1^2) \delta^+((p_1 - P_{in})^2). \quad (2.2)$$

The two delta-functions for $\int d^4 p_1$ is exactly what one needs for the unitarity phase space integration, thus we can use the spinor integration method to evaluate the cross section efficiently.

2.1 The phase space integration with three out-going particles

When $n = 3$ we have

$$I \sim \prod_{i=1}^3 \int d^4 L_i \delta^+(L_i^2) \delta^4(K - \sum L_i) f(L_1, L_2, L_3)$$

$$= \int d^4 L_1 \delta^+(L_1^2) \int d^4 L_2 \delta^+(L_2^2) \delta^+((K - L_1 - L_2)^2) f(L_1, L_2, K - L_1 - L_2)$$

The integration over L_2 with two delta-functions can be done by the spinor integration method. However, for $\int d^4 L_1$ we have only one delta-function. The same difficulty also arises when $n > 3$: for the integration over each momentum, we need another delta function in order to make use of the spinor integration method. Then the problem is how to rewrite the integration measure in such a form where one more delta-function shows up. The solution of this problem is the application of a Faddeev-Popov like method. We will show how it works in detail soon.

Having the solution for the $n = 3$ case, the solution for arbitrary $n \geq 4$ can be obtained by doing the integration recursively one by one. Keeping $(n - 1)$ -momenta fixed, we can integrate the n -th momentum. After that, we keep $(n - 2)$ -momenta fixed and integrate the $(n - 1)$ -th momentum and so on until $n = 3$.

Now let us do the case $n = 3$ explicitly. The integration over L_2 is trivial. We will denote the result after this integration as $\tilde{f}(L_1)$. Then

$$I \sim \int d^4 L_1 \delta^+(L_1^2) \tilde{f}(L_1) \quad (2.3)$$

which is a typical phase space integration and we rewrite it as

$$I = \int \frac{d^4 \ell}{(2\pi)^3} \delta^+(\ell^2) \tilde{f}(\ell). \quad (2.4)$$

To get the standard form suitable for spinor integration method, we need to insert another δ -function. To do so, let us consider the kinematics in more detail. The total energy-momentum tensor of out-going particles is given by K , which has following properties: $K_0 \geq 0$ for positive energy component and $K^2 \geq 0$. With these in mind, let us consider following expression

$$I_z \equiv \int dz \delta((zK - \ell)^2) = \int dz \delta(z^2 K^2 - z(2K \cdot \ell) + \ell^2) = \int dz \sum_{i=1,2} \frac{\delta(z - z_i)}{|2z_i K^2 - (2K \cdot \ell)|}, \quad (2.5)$$

where

$$z_i = \frac{(2K \cdot \ell) \pm \sqrt{(2K \cdot \ell)^2 - 4K^2 \ell^2}}{2K^2}. \quad (2.6)$$

Let's determine the integration region of z for the physical system. In the center-of-mass frame, $K = (E, 0, 0, 0)$, the out-going momentum $\ell = (E_1, P, 0, 0)$ and $E_1^2 = P^2$. Conservation laws ensure that the energy-momentum of all other out-going particles to be $K_{other} = (E - E_1, -P, 0, 0)$. For physical particles we have $K_{other}^2 \geq 0$, i.e., $(E - E_1)^2 \geq P^2$, or $E \geq E_1 + |P|$. Then we get

$$z_i = \frac{2E(E_1 \pm P)}{2E^2} = \frac{E_1 \pm P}{E}. \quad (2.7)$$

When $m = 0$, we have $E_1 = |P|$ and especially $z_- = 0$. Note that $z = 0$ is always a solution when we combine the massless condition $\ell^2 = 0$ in (2.5). Thus to have a physical meaningful solution, we can keep only z_+ solution. It is easy to see that

$$0 \leq z_+ \leq 1 \quad . \quad (2.8)$$

That is, I_z defined in (2.5) makes sense physically only in region $z \in [0, 1]$. With this consideration, I_z can be defined

$$I_z = \int_0^1 dz \frac{\delta(z - z_+)}{|\sqrt{(2K \cdot \ell)^2 - 4K^2 \ell^2}|} = \frac{1}{|\sqrt{(2K \cdot \ell)^2 - 4K^2 \ell^2}|}. \quad (2.9)$$

In another word we have

$$|\sqrt{(2K \cdot \ell)^2 - 4K^2 \ell^2}| \int_0^1 dz \delta((zK - \ell)^2) = 1. \quad (2.10)$$

In the same spirit of the Faddeev-Popov method in gauge-fixings, we insert it into eq (2.3)

$$\begin{aligned} I &= \frac{1}{(2\pi)^3} \int d^4 \ell \delta^+(\ell^2) |\sqrt{(2K \cdot \ell)^2 - 4K^2 \ell^2}| \int_0^1 dz \delta((zK - \ell)^2) \tilde{f}(\ell) \\ &= \frac{1}{(2\pi)^3} \int_0^1 dz \int d^4 \ell \delta^+(\ell^2) (2K \cdot \ell) \delta((zK - \ell)^2) \tilde{f}(\ell). \end{aligned}$$

By use of eq (1.6) and taking the QCD convention $2a \cdot b = \langle a|b|a \rangle$, then

$$\begin{aligned} I &= \frac{-c}{(2\pi)^3} \int_0^1 dz \int \langle \lambda|d\lambda \rangle [\tilde{\lambda}|d\tilde{\lambda}] \int dt dz K^2 \delta \left(z^2 K^2 - zt \langle \lambda|K|\tilde{\lambda} \rangle \right) \tilde{f}(\lambda, \tilde{\lambda}, t) \\ &= \frac{-c}{(2\pi)^3} \int_0^1 dz z \int \langle \lambda|d\lambda \rangle [\tilde{\lambda}|d\tilde{\lambda}] \int dt \delta \left(\frac{zK^2}{\langle \lambda|K|\tilde{\lambda} \rangle} - t \right) \left(\frac{K^2}{\langle \lambda|K|\tilde{\lambda} \rangle} \right)^2 \tilde{f}(\lambda, \tilde{\lambda}, t). \end{aligned}$$

Putting all together we have

$$\begin{aligned} I &= \int \frac{d^4 \ell}{(2\pi)^3} \delta^+(\ell^2) \tilde{f}(\ell) \\ &= \frac{-c}{(2\pi)^3} \int_0^1 dz z \int \langle \lambda|d\lambda \rangle [\tilde{\lambda}|d\tilde{\lambda}] \int dt \delta \left(\frac{zK^2}{\langle \lambda|K|\tilde{\lambda} \rangle} - t \right) \left(\frac{K^2}{\langle \lambda|K|\tilde{\lambda} \rangle} \right)^2 \tilde{f}(\lambda, \tilde{\lambda}, t). \quad (2.11) \end{aligned}$$

where $c = \pi/2$ is related to the Jacobian of changing integration variables and the way we have taken the residues, in which we will omit the $2\pi i$ factor. When the integrand is of the form in $\mathcal{I}_A$,

eq. (2.11) is all one needs to evaluate. When the integrand is of the form in $\mathcal{I}_B$, one needs one more Feynman parametrization to put it in the form of $\mathcal{I}_A$, as discussed in section 1.1.

Now we make two remarks. The first is about the choice of momentum K . It is not arbitrary: the K must be the total energy-momentum tensor of all would-be-integrated momenta. The second is the integration region $z \in [0, 1]$. In fact, the discussion of region of z is nothing, but the discussion of integration boundary faced by the evaluation of cross section. The particular simple result $[0, 1]$ is special for massless particles in 4D. If it is massive, the region of z will be function of mass and total energy-momentum tensor K .

2.2 Arbitrary number of out-going particles and the recursive method

Having worked out the processes of three out-going particles, we can do the integration for arbitrary n out-going particles recursively

$$\begin{aligned} I_n(f; K; P_j) &\equiv \frac{1}{(2\pi)^{3n}} \int \frac{d^3 L_n}{2E_n} \int \frac{d^3 L_{n-1}}{2E_{n-1}} \cdots \int \frac{d^3 L_1}{2E_1} (2\pi)^4 \delta^4(K - \sum_{i=1}^n L_i) f(L_n, L_{n-1}, \dots, L_1) \\ &= \frac{1}{(2\pi)^3} \int \frac{d^3 L_n}{2E_n} I_{n-1}(f; K - L_n; L_n, P_j), \quad (n \geq 3). \end{aligned} \quad (2.12)$$

where in the second line the idea is showing explicitly. More accurately, assuming f is the function of n should-be-integrated momenta L_i , as well as the total energy-momentum tensor $K = \sum_{i=1}^n L_i$ and other external momenta P_j , the integration can be done in two steps. At the first step, we leave the L_n un-integrated, i.e., we integrate other $(n-1)$ momenta L_i . For this case, the total energy-momentum tensor should be $(K - L_n)$. After the first step, we integrate the left L_n again using (2.11) but now with the total energy-momentum tensor K . Again, there is no need to carve out a complicated integration region in the phase space for massless particles. The integrations are always simply over the interval $[0, 1]$.

The idea of recursive evaluation is very natural and has been applied in other methods, for example, straightforward evaluation using momentum components. This direct evaluation will be simplified if we choose right reference frame, for example, the center-of-mass frame. In such case, it usually takes efforts to rewrite results after integration as the Lorentz invariant form. This becomes more severe when we have multiple out-going particles.

In contrast, when the spinor integration method is used recursively, Lorentz invariant expressions are obtained automatically at each step. There is no need to specify any frame and this simplifies the calculation process greatly.

Finally we want to make some observations. First, for each $\int d^3 p$ integration, we have one $\int dz$ integration, thus for n out-going particles, we have $(n-2)$ z -integrations (there maybe other

Feynman parameter integrations from spinor integrations). When $(n - 2)$ is large, usually we can not find the analytic expressions, but since for each z the integration region is $[0, 1]$, the numerical evaluation should be easy to realize.

Secondly, When converting from momentum variables to spinor variables in all integrations, there will be Jacobi factors floating around. In addition, when taking the residues of poles, we have omitted the $2\pi i$ factor. That is to say, normalization factors will be needed in the conversion. In eq. (2.11), it is $c = \pi/2$. Fortunately, for given number of out-going particles, this normalization factor is universal. The simplest way to fix these normalization factors is to calculate the pure physical phase space volume by both methods, as to be shown in the following examples.

3. Example one: two out-going particles

Let us start with the simplest case of two out-going particles. As we have mentioned in introduction, it has exactly the suitable form for the spinor integration method. In the following three subsections, we will use the spinor method to perform the integration. A brief review of method has been presented in the introduction and more details can be found in references [7]-[12]. For comparison, we also list the results from standard momentum integration. In the final subsection, we will briefly discuss the IR/collinear divergences.

As we have mentioned in the introduction, although these examples are very simple and bear little practical importance, they are good for the demonstration of our method, from which the character of our method is clear. The validity of our method for general input shows in the Appendix B.

3.1 The pure phase space integration

We denote the physical phase space integration of n out-going particles as $I_n^{s \text{ or } m}(f; K)$, where s stands for the spinor method and m the momentum method. The K is the sum of momenta of these n particles and f can be a function of K and other external variables as well as the n would-be-integrated momenta.

Spinor integration method : The integration is given by

$$\begin{aligned} I_2^s(1; K) &= \int \frac{d^4 L_2}{(2\pi)^3} \frac{d^4 L_1}{(2\pi)^3} \delta^+(L_2^2) \delta^+(L_1^2) (2\pi)^4 \delta^4(K - L_2 - L_1). \\ &= \frac{-\tilde{c}}{(2\pi)^2} \int t dt \int \langle \lambda | d\lambda \rangle \left[\tilde{\lambda} | d\tilde{\lambda} \right] \delta((K - L)^2) \end{aligned}$$

where $\tilde{c} = \pi/2$ is the Jacobi factor of changing integration variables. As mentioned at the end of the last section, we will omit the $2\pi i$ factor when taking the residues of poles. It has been calculated

in various references [7] and the result is

$$I_2^s(1; K) = \frac{1}{(2\pi)^2} \frac{\pi}{2}. \quad (3.1)$$

Momentum integration method: The expression is given by

$$I_2^m(1; K) = \prod_{i=1,2} \int \frac{dL_i^3}{(2\pi)^3 2E_i} (2\pi)^4 \delta^4(K - L_2 - L_1). \quad (3.2)$$

Taking the center-of-mass frame, where $K = (E, 0, 0, 0)$, and by use of

$$\frac{\delta(E - E_1 - E_2)}{2E_1} = \delta((E - E_2)^2 - E_1^2) = \delta((K - L_2 - L_1)^2) = \delta(E^2 - 2EE_2).$$

we get

$$I_2^m(1; K) = \frac{1}{(2\pi)^2} \int d\Omega \int \frac{E_2^2 dE_2}{2E_2} \delta(E^2 - 2EE_2) = \frac{1}{(2\pi)^2} \frac{\pi}{2}, \quad (3.3)$$

which is identical to the result obtained from the spinor method. It is worth to notice that a suitable choice of reference frame has been made to simplify the calculation.

3.2 The example with $f = 2L_1 \cdot L_2$

Spinor integration method: With this integrand, we have⁷

$$\begin{aligned} I_2^s(u; K) &= \int \frac{d^4 L_2}{(2\pi)^3} \frac{d^4 L_1}{(2\pi)^3} \delta^+(L_2^2) \delta^+(L_1^2) (2\pi)^4 \delta^4(K - L_2 - L_1) 2L_1 L_2 \\ &= \frac{1}{(2\pi)^2} \int d^4 L_2 \delta^+(L_2^2) \delta^+((K - L_2)^2) 2L_2(K - L_2). \end{aligned}$$

Using $L_2^2 = 0$ there is only one term left. Following the process in the first section, we use (1.6) to rewrite the integration and integrate t with delta function, then use (1.15) to get the final result

$$I_2^s(u; K) = \frac{1}{(2\pi)^2} \frac{\pi}{2} K^2. \quad (3.4)$$

Momentum integration method: Similar to the pure phase space integration, we have

$$I_2^m(u; K) = \frac{1}{(2\pi)^2} \frac{\pi}{2} E^2 = \frac{1}{(2\pi)^2} \frac{\pi}{2} K^2 \quad (3.5)$$

which is identical to $I_2^s(u; K)$.

⁷In fact, using $2L_1 \cdot L_2 = (L_1 + L_2)^2 = K^2$, this example is same as the pure phase space integration. Here we do it using different way to demonstrate the technique of spinor integration.

3.3 The example with $f = 1/(2P_1 \cdot L_2 + m^2)$

This example is actually of certain practical value. In real cross section calculations, we meet infrared/collinear divergences when there are massless particles. One way to regularize such divergences is to add a mass term in the propagator.

Spinor integration method: The integral is

$$I_2^s(f; K) = \frac{1}{(2\pi)^2} \int d^4 L_2 \delta^+(L_2^2) \delta^+((K - L_2)^2) \frac{1}{2P_1 \cdot L_2 + m^2}.$$

Using (1.6) to rewrite the integration and integrating t with delta function, we have

$$I_2^s(f; K) = \frac{-1}{(2\pi)^2} \frac{\pi}{2} \int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \frac{1}{\langle \lambda | K | \tilde{\lambda} \rangle \langle \lambda | P_1 + \alpha^2 K | \tilde{\lambda} \rangle}, \quad (3.6)$$

where $\alpha^2 = m^2/E^2$ and the normalization factor $\pi/2$ has been inserted with the spinor integration variables. Introducing a Feynman parameter, we rewrite the above formula as

$$\frac{-1}{(2\pi)^2} \frac{\pi}{2} \int_0^1 dx \int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \frac{1}{\langle \lambda | (1-x)K + x(P_1 + \alpha^2 K) | \tilde{\lambda} \rangle^2}. \quad (3.7)$$

The result can be written down directly

$$I_2^s(f; K) = \frac{1}{(2\pi)^2} \frac{\pi}{2} \int_0^1 dx \frac{1}{((1-x)K + x(P_1 + \alpha^2 K))^2} = \frac{\pi}{2} \frac{1}{(2\pi)^2} \frac{1}{K^2} \ln(1 + \frac{1}{\alpha^2}). \quad (3.8)$$

Momentum integration method: Choosing the center-of-mass frame with $K = P_1 + P_2 = L_1 + L_2$, we have following components $K = (E, 0, 0, 0)$, $P_1 = (E/2, 0, 0, P)$, $P_2 = (E/2, 0, 0, -P)$ and $L_1 = (E_1, -\vec{k}_1)$, $L_2 = (E_2, \vec{k}_1)$, thus $2P_1 L_2 = EE_2(1 - \cos \theta)$, θ is the angle between P_1 and L_2 . Doing the integration we are left with

$$I_2^m(f; K) = \frac{1}{8(2\pi)} \int_0^\pi \sin \theta d\theta \frac{1}{\frac{E^2}{2}(1 - \cos \theta) + m^2} = \frac{1}{(2\pi)} \frac{1}{4E^2} \ln(1 + \frac{1}{\alpha^2}). \quad (3.9)$$

Notice that the result has a logarithm-dependence divergence when $m \rightarrow 0$, which is the usual IR divergence related to massless particles. Again in this calculation, special reference frame has been chosen while in spinor method, each middle step is Lorentz invariant.

3.4 The divergent behavior

To discuss further the divergent behaviors of cross-sections for massless particles, we now consider a simple physical process, in which two gluons are scattered another two $P_1 + P_2 \rightarrow P_3 + P_4$ of different helicity configurations. We will focus on the divergent behaviors in these calculations.

Given color structure, there are three amplitudes with different helicity configurations. Two of them are

$$A_{12}(1^-2^-3^+4^+) = \frac{\langle 1\ 2 \rangle^3}{\langle 2\ 3 \rangle \langle 3\ 4 \rangle \langle 4\ 1 \rangle}, \quad A_{13}(1^-2^+3^-4^+) = \frac{\langle 1\ 3 \rangle^4}{\langle 1\ 2 \rangle \langle 2\ 3 \rangle \langle 3\ 4 \rangle \langle 4\ 1 \rangle}. \quad (3.10)$$

First consider A_{12} and denote $S_{ij} = (P_i + P_j)^2$. In the QCD convention we have⁸

$$f_{12} = |A_{12}|^2 = \frac{S_{12}^2}{S_{23}^2} \quad (3.11)$$

By the momentum method, we have

$$I_2^m(f_{12}; K) = \int \frac{d^3 P_3}{(2\pi)^3} \frac{d^3 P_4}{(2\pi)^3} \frac{1}{2E_3} \frac{1}{2E_4} (2\pi)^4 \delta^4(K - P_3 - P_4) \frac{(2P_1 P_2)^2}{(2P_2 P_3)^2}.$$

Taking the center-of-mass frame and setting the z axis along the direction of $\vec{p}_1$, we have $P_1 = (E, 0, 0, P)$, $P_2 = (E, 0, 0, -P)$, thus the integration is reduced to

$$I_2^m(f_{12}; K) = \frac{1}{2\pi} \int_0^\pi \frac{\sin \theta d\theta}{2(1 - \cos \theta)^2}. \quad (3.12)$$

A singularity appears when $\vec{p}_3$ is parallel to $\vec{p}_2$ and $\theta = 0$. This is the familiar collinear singularity.

The same divergent behavior can be observed in the spinor method. After some elementary algebra, we get the following integration form

$$I_2^s(f_{12}; K) = -\frac{1}{(2\pi)^2} \frac{\pi}{2} \int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \frac{S_{12}}{\langle \lambda | P_2 | \tilde{\lambda} \rangle^2} \sim \frac{1}{P_2^2}. \quad (3.13)$$

Just as we have seen from the momentum method, a divergence appears, since $P_2^2 = 0$.

The situation for A_{13} is a little more complex. We have

$$f_{13} = |A_{13}|^2 = \frac{S_{13}^4}{S_{12}^2 S_{23}^2}.$$

The momentum method gives

$$I_2^m(f_{13}; K) = \frac{1}{2\pi} \int_0^\pi \frac{1}{32} \frac{(1 + \cos \theta)^4}{(1 - \cos \theta)^2} \sin \theta d\theta.$$

It can be rewritten as

$$I_2^m(f_{13}; K) = \frac{1}{2\pi} \int_0^\pi \sin \theta d\theta \left(\frac{1}{2(1 - \cos \theta)^2} - \frac{1}{1 - \cos \theta} \right) + \frac{3}{2(2\pi)} - \frac{1}{2(2\pi)} + \frac{1}{12(2\pi)}. \quad (3.14)$$

⁸It is just one term of the real cross section since we have not added up all color-ordered contribution.

where we have left the first two infinite terms un-integrated.

Now take the spinor method. One has

$$\begin{aligned}
I_2^s(f_{13}; K) &= -\frac{1}{(2\pi)^2} \int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \frac{S_{12} \langle \lambda | 1 | \tilde{\lambda} \rangle^4}{\langle \lambda | 2 | \tilde{\lambda} \rangle^2 \langle \lambda | K | \tilde{\lambda} \rangle^4} \\
&= -\frac{1}{(2\pi)^2} \frac{\pi}{2} \int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \left(\frac{S_{12}}{\langle \lambda | 2 | \tilde{\lambda} \rangle^2} - \frac{4S_{12}}{\langle \lambda | 2 | \tilde{\lambda} \rangle \langle \lambda | K | \tilde{\lambda} \rangle} \right. \\
&\quad \left. + \frac{6S_{12}}{\langle \lambda | K | \tilde{\lambda} \rangle^2} - \frac{4S_{12} \langle \lambda | 2 | \tilde{\lambda} \rangle}{\langle \lambda | K | \tilde{\lambda} \rangle^3} + \frac{S_{12} \langle \lambda | 2 | \tilde{\lambda} \rangle^2}{\langle \lambda | K | \tilde{\lambda} \rangle^4} \right). \tag{3.15}
\end{aligned}$$

where we have used the spinor algebra to split the first line into the second line.

It is not difficult to see that the five terms in $I_2^s(f_{13}; K)$ and $I_2^m(f_{13}; K)$ correspond to each other exactly. In particular, the divergence of $1/(1 - \cos \theta)$ is related to the divergence of $1/\langle \lambda | 2 | \tilde{\lambda} \rangle$.

4. Example two: three out-going particles

4.1 The pure phase space integration

Let us start from the pure phase space integration and identify the normalization factor $c = \pi/2$, as mentioned in section 2.1. From eqs (2.3.20) and (2.3.38) of [13], the phase space integration is reduced to

$$I_3^m(1; K) = \frac{Q^2}{16(2\pi)^3} \int_0^1 dx_1 \int_{1-x_1}^1 dx_2 = \frac{Q^2}{32(2\pi)^3}. \tag{4.1}$$

where $K^2 = Q^2$. In this formula, Lorentz invariant variables x_1, x_2 has been used by initial simplification. This is possible because for small number of particles, there are only a few Lorentz invariant quantities we can construct. With more and more particles, the number of such quantities will increase dramatically.

To make use of the spinor method, we have the integration

$$I_3^s(1; K) = \int \frac{d^4 L_1}{(2\pi)^3} \frac{d^4 L_2}{(2\pi)^3} \frac{d^4 L_3}{(2\pi)^3} \delta^+(L_1^2) \delta^+(L_2^2) \delta^+(L_3^2) (2\pi)^4 \delta^4(K - L_1 - L_2 - L_3). \tag{4.2}$$

The $\int d^4 L_2 d^4 L_3 \delta^+(L_2^2) \delta^+(L_3^2) (2\pi)^4 \delta^4((K - L_1) - L_2 - L_3)$ is just the form of the phase space integration of two outgoing particles, which we know the result is $1/8\pi$. Putting it back we are left

only with

$$\begin{aligned} \frac{\pi}{2} \int \frac{d^4 L_1}{(2\pi)^5} \delta^+(L_1^2) &= \frac{\pi}{2} \frac{-c}{(2\pi)^5} \int_0^1 dz z \int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \int dt \delta \left(\frac{zK^2}{\langle \lambda | K | \tilde{\lambda} \rangle} - t \right) \left(\frac{K^2}{\langle \lambda | K | \tilde{\lambda} \rangle} \right)^2 \\ &= \frac{\pi}{2} \frac{cK^2}{2(2\pi)^5}, \end{aligned} \quad (4.3)$$

Comparing both results we find immediately $c = \pi/2$.

4.2 The example with $f = (L_1 + L_3)^2$

First take the spinor method. Notice that $s = (L_1 + L_3)^2 = (K - L_2)^2$, which can be used to simplify the calculation significantly, since the integrations over L_1 and L_3 are the same as that of the pure phase space integration. Then we have

$$\begin{aligned} I_3^s(s; K) &= \int \frac{d^4 L_2}{(2\pi)^3} \delta^+(L_2^2) (K^2 - 2KL_2) \cdot \frac{1}{8\pi} \\ &= \frac{-\pi^2}{2(2\pi)^5} \int_0^1 dz \frac{z(1-z)}{2} \int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \frac{(K^2)^3}{\langle \lambda | K | \tilde{\lambda} \rangle^2} = \frac{\pi^2 (K^2)^2}{24(2\pi)^5}. \end{aligned} \quad (4.4)$$

Similar to equation (4.1), one has $s = (L_1 + L_3)^2 = Q^2(1 - x_2)$ (as given in [13]), thus

$$I_3^m(s; K) = \frac{Q^2}{16(2\pi)^3} \int_0^1 dx_1 \int_{1-x_1}^1 dx_2 Q^2(1 - x_2) = \frac{(Q^2)^2}{96(2\pi)^3}. \quad (4.5)$$

4.3 The example with $f = s^2 t$

The integration is

$$I_3^s(s^2 t; K) = \prod_{i=1,2,3} \int \frac{d^4 L_i}{(2\pi)^3} \delta^+(L_i^2) (2\pi)^4 \delta^4(K - L_3 - L_2 - L_1) (2L_1 L_3)^2 \cdot 2L_2 L_3. \quad (4.6)$$

This calculation is a little more complicated. However, the basic steps are the same as those for $f = s$. First, integrate over L_1 with the delta function of the energy-momentum conservation; then, integrate over L_2 with the remaining delta functions in spinor coordinates; finally, integrate over L_3 according to eq (2.11). The process is generic.

For illustrations, here we write out the main steps

$$\begin{aligned} I_3^s(s^2 t; K) &= \frac{1}{(2\pi)^5} \int d^4 L_3 \delta^+(L_3^2) \int d^4 L_2 \delta^+(L_2^2) \delta^+((K - L_3 - L_2)^2) \times \\ &\quad ((2KL_3)^2 \cdot 2L_2 L_3 - 2(2KL_3)(2L_2 L_3)^2 + (2L_2 L_3)^3) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{(2\pi)^5} \frac{\pi}{2} \int d^4 L_3 \delta^+(L_3^2) \left(\frac{1}{2} - \frac{2}{3} + \frac{1}{4} \right) (2KL_3)^3 \\
&= -\frac{1}{(2\pi)^5} \frac{\pi^2}{24} \int_0^1 dz \frac{z^4}{2} \int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \frac{(K^2)^5}{\langle \lambda | K | \tilde{\lambda} \rangle^2} = \frac{\pi^2 (K^2)^4}{240(2\pi)^5}.
\end{aligned} \tag{4.7}$$

Using the momentum method [13] we have

$$I_3^m(s^2 t; K) = \frac{Q^2}{16(2\pi)^3} \int_0^1 dx_1 \int_{1-x_1}^1 dx_2 (Q^2)^2 (1-x_2)^2 Q^2 (1-x_1) = \frac{(Q^2)^4}{960(2\pi)^3}, \tag{4.8}$$

which is identical to $I_3^s(s^2 t; K)$.

5. Example three: four out-going particles

In this section we show how to deal with multiple out-going particles recursively. To be concrete, we will focus on the two-four process. Generalization to cases with arbitrary n -outgoing particles will be straightforward.

5.1 The pure phase space integration

Using $I_3^s(1; K)$ as obtained in the previous section and the recursion relation, we have

$$\begin{aligned}
I_4^s(1; K) &= \frac{\pi}{(2\pi)^3} \int d^4 L_4 \delta^+(L_4^2) I_3^s(1; K - L_4) = \frac{1}{4(2\pi)^8} \frac{\pi^2}{2} \int d^4 L_4 \delta^+(L_4^2) (K - L_4)^2 \\
&= \frac{1}{4(2\pi)^8} \frac{\pi^2}{2} \left(K^2 \int d^4 L_4 \delta^+(L_4^2) - \int d^4 L_4 \delta^+(L_4^2) 2KL_4 \right).
\end{aligned}$$

Integrating out these two terms separately, one has

$$I_4^s(1; K) = \frac{(K^2)^2}{768(2\pi)^5}. \tag{5.1}$$

which can compare with the calculation using momentum integration method. Other method, like using the Lorentz invariant variables as $x_{ij} \equiv L_i \cdot L_j$, can also be used to do the calculation. However, unlike the situation with only three L_i , thing becomes more complicated.

5.2 Two examples of non-trivial f

The first example is $f_1 = s$, and the second example is $f_2 = (L_1 + L_2)^2 (L_2 + L_3)^2 (L_3 + L_4)^2 (L_4 + L_1)^2 = (2L_1 L_2)(2L_2 L_3)(2L_3 L_4)(2L_4 L_1)$, which is more complicated. Here the integrations will be performed with the spinor method. The momentum method can be used to make comparisons. For f_1 , the

evaluation is trivial and will not be presented. For f_2 , the calculation is a little involved and will be deferred to the Appendix A.

For the $f_1 = (L_1 + L_2)^2$, we have

$$I_4^s(s; K) = \frac{\pi}{(2\pi)^3} \int d^4 L_4 \delta^+(L_4^2) I_3^s(s; K - L_4) = \frac{1}{(2\pi)^3} \frac{\pi^2}{2} \int d^4 L_4 \delta^+(L_4^2) \frac{((K - L_4)^2)^2}{12(2\pi)^5}, \quad (5.2)$$

by using the result for $I_3^s(s; K)$ obtained in the last section. The last integration is easy to do. Following the prescribed procedure, we get

$$I_4^s(s; K) = \frac{(K^2)^3}{4608(2\pi)^5}. \quad (5.3)$$

Now we turn to the function $f_2 = (2L_1 L_2)(2L_2 L_3)(2L_3 L_4)(2L_4 L_1)$. The factor $(2\pi)^{-8}$ and the Jacobi factor will be dropped for simplicity in the following. The first step is almost the same as that of the momentum method (notice for I_2 , the K is in fact the $K - L_3 - L_4$),

$$\begin{aligned} I_2^s(f_2; K) &= \int d^4 L_2 \delta^+(L_2^2) \int d^4 L_1 \delta^+(L_1^2) \delta^4(K - L_2 - L_1) (2L_1 L_2)(2L_2 L_3)(2L_3 L_4)(2L_4 L_1) \\ &= \int d^4 L_2 \delta^+(L_2^2) \delta^+((K - L_2)^2) \\ &\quad \times ((2L_3 L_4)(2K L_4)(2K L_2)(2L_2 L_3) - (2L_3 L_4)(2K L_2)(2L_2 L_3)(2L_2 L_4)). \end{aligned} \quad (5.4)$$

The first term is simple and it is $(K^2)(2L_3 L_4)(2K L_3)(2K L_4)/2$.

The second term is a little bit more complicated. To proceed, we convert the momentum measure into the spinor measure, and perform the t -integral with the δ -function:

$$\begin{aligned} &\int d^4 L_2 \delta^+(L_2^2) \delta^+((K - L_2)^2) (2L_3 L_4)(2K L_2)(2L_2 L_3)(2L_2 L_4) \\ &= (2L_3 L_4) \int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \frac{(K^2)^4 \langle \lambda | L_3 | \tilde{\lambda} \rangle \langle \lambda | L_4 | \tilde{\lambda} \rangle}{\langle \lambda | K | \tilde{\lambda} \rangle^4}. \end{aligned} \quad (5.5)$$

Then, we do an auxiliary integration as suggested in section 1.1:

$$I_{aux} = \int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \frac{(K^2)^4 \langle \lambda | R | \tilde{\lambda} \rangle^2}{\langle \lambda | K | \tilde{\lambda} \rangle^4} \quad (5.6)$$

with $R = xL_3 + yL_4$. From eq. (1.16), one has

$$I_{aux} = \frac{1}{3} ((K^2)(2KR)^2 - (K^2)^2 R^2). \quad (5.7)$$

Next, pick the coefficient of the (2xy)-term in I_{aux} , which is

$$\frac{1}{3}(K^2)(2KL_3)(2KL_4) - \frac{1}{6}(K^2)^2(2L_3L_4). \quad (5.8)$$

Thus, the end result of (5.5) is

$$\frac{1}{3}(K^2)(2KL_3)(2KL_4)(2L_3L_4) - \frac{1}{6}(K^2)^2(2L_3L_4)^2. \quad (5.9)$$

Combining the two terms in (5.4), we have

$$I_2^s(f; K - L_3 - L_4) = \frac{1}{6}(K^2)(2KL_3)(2KL_4)(2L_3L_4) + \frac{1}{6}(K^2)^2(2L_3L_4)^2. \quad (5.10)$$

There is no more difficulty in remaining steps. After some elementary manipulations, we have

$$I_3^s(f_2; K - L_4) = \int d^4L_3 \delta^+(L_3^2) I_2^s(f_2; K - L_3 - L_4) = \frac{7}{4320}(K^2)^3(2KL_4)^2, \quad (5.11)$$

$$I_4^s(f_2; K) = \int d^4L_4 \delta^+(L_4^2) I_3^{s,r}(f_2; K - L_4) = \frac{1}{172800}(K^2)^6. \quad (5.12)$$

Putting all factors back, we get finally

$$I_4^s(f_2; K) = \frac{1}{675 \times 2^{12}(2\pi)^5}(K^2)^6. \quad (5.13)$$

as to be checked with the calculations in the Appendix A.

6. Conclusion

In this paper, we have proposed to use the spinor integration method, which is developed from the phase space integration of one-loop unitarity cut, to do the real physical phase space integration for the total cross section. This paper, as the initial construction of frame, focus on the massless particles in pure 4D. We will discuss the massive and general D-dimension case in subsequent work.

Now, let us sum some salient points of our method. The first point is the rewriting measure given in (2.11) by the Faddeev-Popov trick. This new form enable us to do the spinor integration method. Since the z -integration is always $[0, 1]$ for massless case, we can change the order of integration and leave all z -integration at the end after we have performed all spinor integrations. With n out-going particles, there will be $(n - 2)$ z -integration to do. The integrand will be the Lorentz invariant expression of momenta of in-coming particles only. With two particles p_1, p_2 we have only one nontrivial variables $K^2 = (p_1 + p_2)^2$. For massive case, the integrand for z will depend on K as well as various masses and thing will be more complicated.

In comparison, there will be $(3n-4)$ integrations in general, if we use the momentum integration method⁹. Furthermore, the expressions are usually in component-form which is not manifest Lorentz invariant. With more and more out-going particles, it will also become difficult to specify the integrated regions and separate angles variables and module variables.

One can also try to rewrite the measure using Lorentz invariant variables. With n L_i there are $\frac{n(n-1)}{2}$ quantities given by $x_{ij} = L_i \cdot L_j$. They are not independent to each other in general, so the discussion of proper choice of subset as well as the constraints among them becomes more and more tedious with the increase of number of L_i .

Having shown the promise of our new method, the result in this paper is not immediately to be useful for practical calculations. As we have mentioned several times, we need to include massive outgoing particles as well as general D-dimension. This can, in fact, be accomplished by mimicking one-loop calculations, where one generalizes pure 4D spinor integration to $(4-2\epsilon)$ dimension spinor integration [9, 10] although some technical difficulties need to be attacked. Work on this is in progress.

Acknowledgement: We would like to thank C. Anastasiou, R. Britto, and Z. Kunszt for reading the draft. B.F would also like to thank the hospitality of CUFE where the final part is done. The work is funded by Qiu-Shi funding from Zhejiang University and Chinese NSF funding under contract No.10875104, 10425525 and 10875103.

A. Using momentum method to integrate f_2 for four out-going particles

In this appendix, we will present the integration with $f_2 = (2L_1L_2)(2L_2L_3)(2L_3L_4)(2L_4L_1)$ for four out-going particles by using the momentum method. It is to be compared with the spinor method presented in section 5.2. For simplicity, the factor $(2\pi)^{-8}$ has been dropped, just as in section 5.2. In the spirit of recursion, we first perform the integration

$$\begin{aligned} I_2^m(f_2; K) &= \int \frac{d^3L_2}{2E_2} \int \frac{d^3L_1}{2E_1} \delta^4(K - L_2 - L_1) (2L_1L_2)(2L_2L_3)(2L_3L_4)(2L_4L_1) \\ &= \int \frac{d^3L_2}{2E_2} \delta^+((K - L_2)^2) \\ &\quad \times ((2L_3L_4)(2KL_4)(2KL_2)(2L_2L_3) - (2L_3L_4)(2KL_2)(2L_2L_3)(2L_2L_4)). \end{aligned} \quad (\text{A.1})$$

In the following, K will be substituted by $K - L_3 - L_4$. The first term is simpler than the second. It is

$$\int \frac{d^3L_2}{2E_2} \delta^+((K - L_2)^2) (2L_3L_4)(2KL_4)(2KL_2)(2L_2L_3)$$

⁹With a suitable choice of reference framework, we may reduce dimension of integration somewhat further.

$$\begin{aligned}
&= (2L_3L_4)(2KL_4) \int \frac{d^3L_2}{2E_2} \delta^+(E^2 - 2EE_2)(2EE_2) \left(2E_2E_3(1-z) \right), \quad (z \equiv \cos \theta_{23}) \\
&= (2L_3L_4)(2KL_4) \cdot \frac{2\pi}{8} (E^2)(2EE_3) = \frac{2\pi}{8} (K^2)(2KL_4)(2KL_3)(2L_3L_4). \tag{A.2}
\end{aligned}$$

In the last line, we have transformed from the component-form to the Lorentz-invariant form, such as $2EE_3 \rightarrow 2KL_3$, to make it suitable for the use of recursion relations. Such manipulations will always be used in the following.

Next we deal with the second term (without the minus sign). The difficulty is that the integrand depends on two angles, θ_{23} and θ_{24} , which cannot be integrated over at the same time. However we can spilt the integral into two terms, either of which has only one angle. To do this, we introduce two auxiliary parameters $L_+ = L_3 + L_4$ and $L_- = L_3 - L_4$. Then the second term becomes

$$\begin{aligned}
&\int \frac{d^3L_2}{2E_2} \delta^+((K - L_2)^2)(2L_3L_4)(2KL_2)(L_2(L_+ + L_-))(L_2(L_+ - L_-)) \\
&= (2L_3L_4) \int \frac{d^3L_2}{2E_2} \delta^+((K - L_2)^2)(2KL_2) [(L_2L_+)^2 - (L_2L_-)^2]. \tag{A.3}
\end{aligned}$$

Now we calculate the first term of the expression only. The second term is almost the same as the first. Set $L_+ = (E_+, P_+\hat{p}_+)$, $L_- = (E_-, P_-\hat{p}_-)$, where P_+ and P_- are the module of the corresponding momentum vector. θ_{2+} is the angle between $\vec{l}_2$ and $\hat{p}_+$, and θ_{2-} is that between $\vec{l}_2$ and $\hat{p}_-$. Then the first term becomes

$$\begin{aligned}
&(2L_3L_4) \int_0^{2\pi} d\varphi \int_{-1}^1 dz_+ \int \frac{E_2^2}{2E_2} dE_2 \delta^+(E^2 - 2EE_2)(2EE_2)(E_2E_+ - E_2P_+z_+)^2, \quad (z_+ \equiv \cos \theta_{2+}) \\
&= (2\pi)(2L_3L_4) \left(\frac{E^2}{16} (EE_+)^2 + \frac{E^4}{48} P_+^2 \right). \tag{A.4}
\end{aligned}$$

Also we have $L_+^2 = E_+^2 - P_+^2 = (L_3 + L_4)^2 = 2L_3L_4$, i.e. $P_+^2 = E_+^2 - 2L_3L_4$. Substituting it into eq (A.4), we have

$$(2\pi) \left(\frac{1}{12} (K^2)(2L_3L_4)(KL_+)^2 - \frac{1}{48} (K^2)^2(2L_3L_4)^2 \right).$$

Similarly, the second term gives

$$(2\pi) \left(\frac{1}{12} (K^2)(2L_3L_4)(KL_-)^2 + \frac{1}{48} (K^2)^2(2L_3L_4)^2 \right).$$

Combining them together, we get the result of the second term in (A.1):

$$(2\pi) \left(\frac{1}{12} (K^2)(2KL_4)(2KL_3)(2L_3L_4) - \frac{1}{24} (K^2)^2(2L_3L_4)^2 \right). \tag{A.5}$$

Then

$$\begin{aligned} I_2^m(f_2; K) &= \left(\frac{2\pi}{8} - \frac{2\pi}{12} \right) (K^2)(2KL_4)(2KL_3)(2L_3L_4) + \frac{2\pi}{24}(K^2)^2(2L_3L_4)^2 \\ &= \frac{2\pi}{24}(K^2)(2KL_4)(2KL_3)(2L_3L_4) + \frac{2\pi}{24}(K^2)^2(2L_3L_4)^2. \end{aligned} \quad (\text{A.6})$$

The rest calculations are tedious but not difficult. The end results are

$$I_3^m(f_2; K) = \int \frac{d^3L_3}{2E_3} I_2^m(f_2; K - L_3) = \frac{7}{34560} (2\pi)^2 (K^2)^3 (2KL_4)^2, \quad (\text{A.7})$$

$$I_4^m(f_2; K) = \int \frac{d^3L_4}{2E_4} I_3^m(f_2; K - L_4) = \frac{(2\pi)^3}{2764800} (E^2)^6. \quad (\text{A.8})$$

Adding back the factor of $(2\pi)^{-8}$, we finally get

$$I_4^m(f_2; K) = \frac{1}{2764800(2\pi)^5} (E^2)^6. \quad (\text{A.9})$$

which is the same as $I_4^s(f_2; K)$, as obtained in section 5.2.

B. The spinor integration as integration in complex plane

In this section, we will rewrite spinor integration as the integration in the complex plane. The correctness of the spinor integration method will be obvious. Furthermore, as complex integration can be applied to any form of inputs in principal, so can the spinor integration be. This is very important since for phase space integration, the input has singularities other than those in the propagators in the one-loop calculation (see, for example, [15]). We will demonstrate it with a few examples.

B.1 Rewriting of spinor integration

Since spinor λ has only two components, we can expand it with two independent spinors, for example, λ_a, λ_b as

$$|\lambda\rangle = |a\rangle + z|b\rangle, \quad |\tilde{\lambda}\rangle = |a\rangle + \bar{z}|b\rangle. \quad (\text{B.1})$$

For real momentum, $\bar{z}$ is the complex conjugation of z . Using this expansion, we have $\langle\lambda|d\lambda\rangle = \langle a|b\rangle dz$, $[\tilde{\lambda}|d\lambda] = [a|b]d\bar{z}$, thus the measure is given by

$$\int \langle\lambda|d\lambda\rangle [\tilde{\lambda}|d\lambda] = \langle a|b\rangle [a|b] \int dzd\bar{z}. \quad (\text{B.2})$$

With an arbitrary integrand $G(\lambda, \tilde{\lambda})$, using above replacement we have

$$\int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] G(\lambda, \tilde{\lambda}) = \langle a | b \rangle [a | b] \int dz d\bar{z} G(z, \bar{z}) . \quad (\text{B.3})$$

In another word, the spinor integration is a two-dimensional integration over complex plane.

For integration in complex plane, there is an important formula [14]

$$\oint_C (u dx + v dy) = \int \int_R \left(\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) \quad (\text{B.4})$$

or in the form

$$\oint_C g dz = - \int \int_R \frac{\partial g}{\partial \bar{z}} dz \wedge d\bar{z} . \quad [\text{S-form-1}] \quad (\text{B.5})$$

Now we see how to apply above formula with the simplest example:

$$\begin{aligned} & \int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \frac{1}{\langle \lambda | K | \lambda \rangle^2} = \int \int \langle a | b \rangle [a | b] dz d\bar{z} \frac{1}{(\langle \lambda | K | a \rangle + \bar{z} \langle \lambda | K | b \rangle)^2} \\ &= \int \int \langle a | b \rangle [a | b] dz d\bar{z} \frac{d}{d\bar{z}} \frac{1}{\langle \lambda | K | b \rangle} \frac{-1}{(\langle \lambda | K | a \rangle + \bar{z} \langle \lambda | K | b \rangle)} \\ &= - \oint \langle a | b \rangle [a | b] dz \frac{1}{\langle \lambda | K | b \rangle} \frac{-1}{(\langle \lambda | K | a \rangle + \bar{z} \langle \lambda | K | b \rangle)} . \end{aligned} \quad (\text{B.6})$$

Using $\langle \lambda | K | b \rangle = \langle a | K | b \rangle + z \langle b | K | b \rangle$ with pole location $z = -\frac{\langle a | K | b \rangle}{\langle b | K | b \rangle}$. Taking residue of pole we get $\frac{-1}{K^2}$ (where we have neglected the $2\pi i$ factor).

It will be useful to compare above calculation with the one we did in spinor integration. We have

$$\int \langle \lambda | d\lambda \rangle [\tilde{\lambda} | d\tilde{\lambda}] \frac{1}{\langle \lambda | K | \lambda \rangle^2} = \int \langle \lambda | d\lambda \rangle [d\tilde{\lambda} | \partial_{\tilde{\lambda}}] \frac{[\eta | \tilde{\lambda}]}{\langle \lambda | K | \eta \rangle \langle \lambda | K | \tilde{\lambda} \rangle} . \quad (\text{B.7})$$

Taking the residue of $\langle \lambda | K | \eta \rangle$ we have $\frac{-1}{K^2}$. If we set $\eta = b$ and using

$$d\bar{z} \frac{\partial}{\partial \bar{z}} = [d\tilde{\lambda} | \partial_{\tilde{\lambda}}] , \quad (\text{B.8})$$

we will see explicitly the one-to-one correspondence between our spinor integration method and complex integration method.

The power of (B.5) is that it reduces the two-dimension integration into pure algebraic calculation, i.e., reading out residue by implicitly using

$$\frac{\partial}{\partial \bar{z}} \frac{1}{z - a} = \pi \delta^2(z - a) . \quad (\text{B.9})$$

Spinor integration method has also used this property, which is casted into the spinor formula and called the “holomorphic anomaly” [5].

It is obvious now that any spinor integration can be rewritten as complex plane integration, so its validity is obvious. However, keeping the spinor form the expression will be much more compact and also easier to read out residues.

B.2 More examples

The spinor method is developed for the one-loop calculation at beginning, thus the most cases we have met are propagator-like singularities, where we have result (1.15). For other kinds of singularities, we need similar results. This can be obtained by first going to complex variables and then pulling back to spinor variables. Using this technique we can do the spinor integration for other kinds of singularities. Let us demonstrate this with a few examples.

First let us discuss

$$\begin{aligned}
& \int \langle \lambda \, d\lambda \rangle [\lambda \, d\lambda] \frac{1}{\langle \lambda | K | \lambda \rangle \langle \lambda | R | \lambda \rangle} \\
&= \int \langle a \, b \rangle [a \, b] dz d\bar{z} \frac{1}{(\langle \lambda | K | a \rangle + \bar{z} \langle \lambda | K | b \rangle)(\langle \lambda | R | a \rangle + \bar{z} \langle \lambda | R | b \rangle)} \\
&= \int \langle a \, b \rangle [a \, b] dz d\bar{z} \left(\frac{\langle \lambda | K | b \rangle}{\langle \lambda | K | a \rangle + \bar{z} \langle \lambda | K | b \rangle} - \frac{\langle \lambda | R | b \rangle}{\langle \lambda | R | a \rangle + \bar{z} \langle \lambda | R | b \rangle} \right) \\
&\quad \times \frac{1}{\langle \lambda | K | b \rangle \langle \lambda | R | a \rangle - \langle \lambda | R | b \rangle \langle \lambda | K | a \rangle} \\
&= \int \langle a \, b \rangle d\bar{z} \frac{\partial}{\partial \bar{z}} (\ln(\langle \lambda | K | a \rangle + \bar{z} \langle \lambda | K | b \rangle) - \ln(\langle \lambda | R | a \rangle + \bar{z} \langle \lambda | R | b \rangle)) \\
&\quad \times \frac{1}{\langle \lambda | K R | \lambda \rangle}
\end{aligned} \tag{B.10}$$

Using (B.8) as well as

$$\left[\alpha \left| \frac{\partial}{\partial \tilde{\lambda}} \right. \right] [\beta | \tilde{\lambda}] = [\beta | \alpha] \tag{B.11}$$

above result is equivalent to

$$\left[d\tilde{\lambda} \, \partial_{\tilde{\lambda}} \right] \left(\frac{1}{\langle \lambda | K | R | \lambda \rangle} \ln \frac{\langle \lambda | K | \tilde{\lambda} \rangle}{\langle \lambda | R | \tilde{\lambda} \rangle} \right) = \left[\tilde{\lambda} \, d\tilde{\lambda} \right] \frac{1}{\langle \lambda | K | \tilde{\lambda} \rangle \langle \lambda | R | \tilde{\lambda} \rangle} \tag{B.12}$$

Similarly we have

$$\left[d\tilde{\lambda} \partial_{\tilde{\lambda}} \right] \left(\frac{1}{\langle \lambda | K | R | \lambda \rangle} \sqrt{\frac{\langle \lambda | R | \tilde{\lambda} \rangle}{\langle \lambda | K | \tilde{\lambda} \rangle}} \right) = \frac{-1}{2} \left[\tilde{\lambda} d\tilde{\lambda} \right] \frac{1}{\langle \lambda | K | \tilde{\lambda} \rangle^{\frac{3}{2}} \langle \lambda | R | \tilde{\lambda} \rangle^{\frac{1}{2}}} \quad (\text{B.13})$$

Using these results, we can do more examples:

Example 1: The first example is the square root form as

$$\begin{aligned} I_1 &= \int d^4 L_1 \delta^+(L_1^2) \delta^+((L_1 - K)^2) \frac{1}{\sqrt{L_1 \cdot R}} \\ &= \int \langle \lambda d\lambda \rangle \left[\tilde{\lambda} d\tilde{\lambda} \right] \sqrt{\frac{2}{K^2}} \frac{1}{\langle \lambda | K | \tilde{\lambda} \rangle^{3/2} \sqrt{\langle \lambda | R | \tilde{\lambda} \rangle}} \\ &= \int \langle \lambda d\lambda \rangle \frac{2\sqrt{\frac{2}{K^2}}}{\langle \lambda | K | R | \lambda \rangle} \left[d\tilde{\lambda} \left| \frac{\partial}{\partial \tilde{\lambda}} \right| \right] \frac{-\sqrt{\langle \lambda | R | \tilde{\lambda} \rangle}}{\sqrt{\langle \lambda | K | \tilde{\lambda} \rangle}} \end{aligned} \quad (\text{B.14})$$

Reading out residues we have

$$I_1 = \frac{2(\sqrt{2K \cdot R - \sqrt{\Delta}} + \sqrt{2K \cdot R + \sqrt{\Delta}})}{K^2 \sqrt{\Delta}} \quad (\text{B.15})$$

where $\Delta = (2R \cdot K)^2 - 4R^2 K^2$.

Example 2:

The second example is the one with gram determinant. This is, in fact, the one we will meet in one-loop calculation as well as in tree-level after one external particle having been integrated out. It represents the typical singularity behavior in the phase space integration.

$$\begin{aligned} I_2 &= \int d^4 L \delta^+(L^2) \delta^+((L - K)^2) = \int d^4 L \delta^+(L^2) \delta^+((L - K)^2) \frac{1}{\det \begin{pmatrix} K_1^2 & K_1 \cdot K_2 & K_1 \cdot K_3 \\ K_1 \cdot K_2 & K_2^2 & K_3 \cdot K_2 \\ K_1 \cdot K_3 & K_3 \cdot K_2 & K_3^2 \end{pmatrix}} \\ &= \int \frac{d^4 L \delta^+(L^2) \delta^+((L - K)^2)}{2(K \cdot R)(L \cdot R)(K \cdot L) - R^2(K \cdot L)^2 - K^2(L \cdot R)^2} \end{aligned}$$

where we have taken $K_1 = K, K_2 = R, K_3 = L$. Doing standard manipulation we get

$$\begin{aligned}
I_2 &= \int \langle \lambda \, d\lambda \rangle [\lambda \, d\lambda] \frac{-4}{K^2} \frac{1}{2(K \cdot R) \langle \lambda | K | \lambda \rangle \langle \lambda | R | \lambda \rangle - R^2 \langle \lambda | K | \lambda \rangle^2 - K^2 \langle \lambda | R | \lambda \rangle^2} \\
&= \int \langle \lambda \, d\lambda \rangle [\lambda \, d\lambda] \frac{-4}{K^2} \frac{1}{\langle \ell | x_1 R + y_1 K | \ell \rangle \langle \ell | x_2 R + y_2 K | \ell \rangle} \quad [\text{I2-fac}] \tag{B.16}
\end{aligned}$$

where

$$y_2 = 1, \quad y_1 = -R^2, \quad x_1 = K \cdot R + \sqrt{(K \cdot R)^2 - K^2 R^2}, \quad x_2 = \frac{-K \cdot R + \sqrt{(K \cdot R)^2 - K^2 R^2}}{R^2}$$

Now this is the familiar case given by (1.13) and we have the final result

$$\begin{aligned}
I_2 &= \frac{-4}{K^2} \int_0^1 d\alpha \frac{1}{((1-\alpha)(x_1 R + y_1 K) + \alpha(x_2 R + y_2 K))^2} \\
&= \frac{1}{K^2((K \cdot R)^2 - K^2 R^2)} \int_0^1 d\alpha \frac{1}{\alpha(\alpha - 1)} \tag{B.17}
\end{aligned}$$

which is divergent with α -integration. This example is, in fact, another typical example of divergence at the boundary of phase space.

It is worth to emphasize that the factorization (B.16) we have done is the general method we need to use for phase space integration in more complicated situations.

References

- [1] Z. Bern *et al.* [NLO Multileg Working Group], arXiv:0803.0494 [hep-ph].
- [2] Z. Bern, L. J. Dixon, D. C. Dunbar and D. A. Kosower, Nucl. Phys. B **425**, 217 (1994) [arXiv:hep-ph/9403226].
- [3] Z. Bern, L. J. Dixon, D. C. Dunbar and D. A. Kosower, Nucl. Phys. B **435**, 59 (1995) [arXiv:hep-ph/9409265].
- [4] E. Witten, Commun. Math. Phys. **252**, 189 (2004)
- [5] F. Cachazo, P. Svrcek and E. Witten, JHEP **0410**, 077 (2004)
- [6] R. Britto, F. Cachazo and B. Feng, Phys. Rev. D **71**, 025012 (2005)
- [7] R. Britto, E. Buchbinder, F. Cachazo and B. Feng, Phys. Rev. D **72**, 065012 (2005) [arXiv:hep-ph/0503132].
- [8] R. Britto, B. Feng and P. Mastrolia, Phys. Rev. D **73**, 105004 (2006)

- [9] C. Anastasiou, R. Britto, B. Feng, Z. Kunszt and P. Mastrolia, Phys. Lett. B **645**, 213 (2007) [arXiv:hep-ph/0609191].
- [10] C. Anastasiou, R. Britto, B. Feng, Z. Kunszt and P. Mastrolia, JHEP **0703**, 111 (2007) [arXiv:hep-ph/0612277].
- [11] A. Gehrmann-De Ridder, T. Gehrmann and G. Heinrich, Nucl. Phys. B **682**, 265 (2004) [arXiv:hep-ph/0311276].
- [12] R. Britto and B. Feng, Phys. Rev. D **75**, 105006 (2007) [arXiv:hep-ph/0612089].
- [13] R.D. Field, "Application of Perturbative QCD", Addison-Wesley Publishing Company, 1989.
- [14] M. J. Ablowitz and A. S. Fokas, "Complex Variables: Introduction and Applications", Cambridge University Press, 1997.
- [15] M. Czakon and A. Mitov, Nucl. Phys. B **824**, 111 (2010) [arXiv:0811.4119 [hep-ph]].